

Published in final edited form as:

J Med Chem. 2012 October 25; 55(20): 8582–8587. doi:10.1021/jm300612z.

(S)-N-(2,5-Dimethylphenyl)-1-(quinoline-8-ylsulfonyl)pyrrolidine-2-carboxamide as a Small Molecule Inhibitor Probe for the Study of Respiratory Syncytial Virus Infection

Blake P. Moore^{†,‡}, Dong Hoon Chung[†], Daljit S. Matharu[†], Jennifer E. Golden[†], Clinton Maddox[†], Lynn Rasmussen[†], James W. Noah[†], Melinda I. Sosa[†], Subramaniam Ananthan[†], Nichole, A. Tower[†], E. Lucile White[†], Fuli Jia^{†,§}, Thomas E. Prisinzano^{†,§}, Jeffrey Aubé^{†,§}, Colleen B. Jonsson[⊥], and William E. Severson^{*,⊥}

[†]Southern Research Specialized Biocontainment Screening Center, Southern Research Institute, Birmingham, AL, USA

[‡]University of Kansas Specialized Chemistry Center, University of Kansas, Lawrence, KS, USA

[§]Department of Medicinal Chemistry, University of Kansas, Lawrence, KS, USA

[⊥]Center for Predictive Medicine for Biodefense and Emerging Infectious Diseases, University of Louisville, Louisville, KY, USA

Abstract

A high-throughput, cell-based screen was used to identify chemotypes as inhibitors for human respiratory syncytial virus (*hRSV*). Optimization of a sulfonylpyrrolidine scaffold resulted in compound **5o** that inhibited a virus-induced cytopathic effect in the entry stage of infection ($EC_{50} = 2.3 \pm 0.8 \mu M$) with marginal cytotoxicity ($CC_{50} = 30.9 \pm 1.1 \mu M$) and reduced viral titer by 100-fold. Compared to ribavirin, sulfonylpyrrolidine **5o** demonstrated an improved *in vitro* potency and selectivity index.

INTRODUCTION

Human respiratory syncytial virus (*hRSV*) is the most common cause of bronchiolitis and pneumonia among infants and children under one year of age.¹ In the United States there are approximately 125,000 yearly *hRSV*-related hospitalizations, and of those, 500 young children could die due to the infection or its complications each year.² The virus is highly contagious and affects those with compromised cardiac, pulmonary (COPD), and immune systems.³ As such, the elderly are also a highly susceptible population to *hRSV*. Treatment options are limited. Due to infant mortality associated with attempted vaccination, vaccine development is proceeding cautiously.^{4,5} Synagis® (Palivizumab), a humanized monoclonal antibody, is a prophylactic, injectable therapeutic used only with high risk pediatric patients.⁶ Ribavirin, a nucleoside antimetabolite, is approved for acute infection^{7,8} and infected, immunocompromised patients,⁹ but has a long half-life and accumulates in erythrocytes, thus requiring regeneration of the affected red blood cells to eliminate the

*Corresponding Author Phone (502) 852-1546, weseve01@louisville.edu.

[‡]Current address: Department of Biochemistry & Molecular Genetics, University of Alabama, Birmingham, AL, USA

[§]Current Address: Department of Molecular & Cellular Biology, Baylor College of Medicine, Houston, TX, USA

Supporting Information. Assay details and experimental characterization for select compounds. This material is available free of charge via the internet at <http://pubs.acs.org>.

drug. These issues, coupled with embryocidal and teratogenic effects constitute severe toxicological liabilities that limit the use of ribavirin, especially in infants,⁷ around administering pregnant medical personnel, and in treated partners of pregnant women (Figure 1).

To address the absence of clinically relevant and safe *h*RSV therapies, many investigators have pursued a target-based drug design approach. Ribavirin 5'-monophosphate resembles GMP and can decrease cellular GTP pools due to the inhibition of the enzyme inosine monophosphate dehydrogenase (IMPDH).^{10,11} Nevertheless, this decrease does not completely account for the observed antiviral activity as inhibitory effects have been noted on RNA capping¹² and direct inhibition of viral polymerase activity for influenza viruses.¹³ Literature disclosed IMPDH inhibitors have reported *in vitro* therapeutic indices that are not competitive with ribavirin.¹⁴ Several other inhibitors that target the fusion protein,^{15–20} ribonucleoprotein (RNP) complex,²¹ guanylation events,²² and N-protein²³ have been discovered. Of these, some demonstrate potency and limited toxicity in animal models.^{24–26} However, due to formulation for oral bioavailability,¹⁹ strategic reasons,²⁷ and observed loss of activity *in vivo*,²⁸ many have not progressed to the clinic to combat *h*RSV disease.

In an effort to identify lead compounds acting through new mechanisms, we developed, optimized and validated a high-throughput cell-based screen that measures the respiratory syncytial virus-induced cytopathic effect (CPE) in HEp-2 cells (unpublished results). CPE was measured using the Cell Titer-Glo™ viability assay in which the luminescent signal generated is directly proportional to the amount of cellular ATP present which is also proportional to the number of metabolically active cells. A total of 313,816 compounds from the Molecular Libraries Small Molecule Repository (MLSMR) were screened in single dose against *h*RSV (strain Long) at a concentration of 10 μ M. Hits (2,465 compounds) were evaluated for their antiviral activity and cell toxicity in dose response experiments, and 409 compounds produced a protective effect of at least 50% CPE inhibition. Based on potency, selectivity and chemical tractability, 51 hits were selected for verification in an *in vitro* titer reduction assay to assess their effect on the production of infectious progeny virus. Many chemotypes of interest emerged, including several compounds containing a sulfonylpyrrolidine moiety. Hit compound **2** displayed a CPE EC₅₀ of 5.0 μ M, a CC₅₀ (HEp-2 cellular cytotoxicity) of 31.5 μ M, and a selectivity index (CC₅₀/EC₅₀) of 6.3 (Figure 2). Profile improvements were explored by modulating various structural elements (shaded regions, Figure 2).

CHEMISTRY

Analogs were prepared by treating substituted sulfonyl chlorides **3** with proline or another suitable amino acid to generate sulfonamide carboxylates **4**. Coupling of the acids to various amines via traditional methods afforded the desired products **5** (Scheme 1).

RESULTS AND DISCUSSION

Attention was first placed on the proline unit between the aryl sulfone and aniline groups. The hit compound **2** was tested as a racemic mixture; therefore, each enantiomer was individually prepared from *L*- or *D*-proline to determine the more pharmacologically active constituent. The *S*-enantiomer, derived from *L*-proline, was found to be the more active component of the racemic mixture, delivering a selectivity index of 11.8 when cytotoxicity was accounted for (entry 4, Table 1). This compound showed a 1 log reduction in virus titer in a plaque reduction assay. Acyclic variants of the linker region, including those that probed the methylation of the amide nitrogen, were found to have EC₅₀ values > 50.0 μ M.

Several variants of the quinoline moiety were pursued but offered no benefit in potency. Replacing the nitrogen atom with –CH– afforded an inactive analog ($EC_{50} > 50.0 \mu M$), as did migrating the nitrogen to alternate positions of the quinoline substructure. The replacement of the quinoline with a 4-linked-benzoxadiazole, a phenyl ring or a simple methyl group was also not advantageous. Substitution of the sulfonyl group (SO_2) for a carbonyl functionality or its replacement with –CH₂– resulted in complete loss of potency ($EC_{50} > 50.0 \mu M$, see supporting information, Tables S1 and S2, respectively). Focus was then shifted to the modification of the 2,4-dimethylanilide. Following the results noted above, all analogs in this study were prepared as the *S*-enantiomer (Table 2). Simplifying the 2,4-dimethylphenyl substitution pattern down to monomethyl substituted phenyl derivatives revealed that the observed potency was not due to the presence of one group alone (entries 3–5). Increased steric bulk at C2 marginally improved potency vs. monomethyl substitution at the same position (*cf.* entries 8 to 7). Mimicking the 2,4-substitution pattern with chlorine atoms in place of methyl groups resulted in complete loss of potency (**5n**, entry 11). As the 2-alkyl substituent appeared to be necessary in combination with other substituents to preserve potency, this dynamic was explored further to reveal that the 2,5-dimethylphenyl moiety of analog **5o** was slightly more beneficial in terms of potency and maintained the cytotoxicity threshold (entry 12, $EC_{50} = 2.3 \mu M$, $CC_{50} = 30.9 \mu M$). For select compounds assessed for aqueous solubility, no effect was observed on CPE potency. Solubility and stability were determined for **5o**, revealing an acceptable solubility measurement of 92.7 $\mu g/mL$ in PBS buffer and stability of 95.4% (unchanged parent remaining) after 48 h in 50% PBS/50% acetonitrile.²⁹

To probe the mechanism of action of the sulfonylpyrrolidines, the window of inhibitory activity in the cell-based assay was refined. Potency of compounds over time following infection was examined to ascertain early (entry) or late (replication) antiviral activity in the virus life cycle.³⁰

In the time of addition study, HEp-2 cells were infected with *h*RSV strain Long at an MOI of 3.0 at time point 0 and incubated for 6 days and test compounds, **5b**, **5t**, **5o**, or ribavirin were added to plates at –1, 0, 1, 2, 3, 5, 7, 21 and 24 h post infection (p.i.). CPE was assessed using Cell-Titer Glo as an endpoint reagent. Controls without test compound included HEp-2 cells with no *h*RSV exposure (cell control) and *h*RSV-infected cells (virus control). To evaluate cellular toxicity attributed to test compound alone, uninfected HEp-2 cells were treated with **5b**, **5t**, **5o**, or ribavirin at 25 μM concentration at time point 0, and cell viability was assessed after 144 hours.

There was less than 1% cell viability for the *h*RSV-infected cells without addition of any test compound or ribavirin. Uninfected cells treated with ribavirin, **5t** or **5o** displayed 90%, 88% and 95% cell viability, respectively, indicating low cellular toxicity inherent to these compounds. However, uninfected cells treated with **5b** exhibited only 46% cellular viability, suggesting moderate toxicity due to the test compound alone.³¹ Ribavirin treatment protected cells from *h*RSV induced CPE for up to 7 h p.i., indicating that it targets the period of infection during which viral replication is in progress. Compound **5o** protected cells from *h*RSV induced CPE (> 50%) from 1 – 3 h p.i., and at 24 h p.i. cell viability was only 26%. (Figure 3).

Two analogs of similar profile to **5o**, **5b** and **5t**, demonstrated a decrease in efficacy when added at each time point from 0–5 hours p.i. This profile could be due to the inhibition of one or more early virus life cycle steps (entry, post-entry, or early-stage infection processes), a hypothesis that is supported by an inability of these compounds to affect processes later in the viral replication cycle. This data lead us to conclude that **5o** was inhibiting early infection events, characterized by viral attachment, uptake, fusion or initial transcription.

The sulfonylpyrrolidine scaffold analogues were evaluated for their ability to reduce the amount of infectious virus produced in cell culture. These measurements of compound-mediated viral titer reduction were used to complement the cytoprotection assay results. A standard plaque-reduction assay was used as a secondary assay to determine the ability of this class of compounds to reduce the amount of infectious virus produced in HEp-2 cells.³² Cells were infected with *h*RSV in the presence of 25 μ M test compound (**5b**, **5o**, or **5t**). Compounds **5b** and **5t** each showed 1 log reduction in virus titer, or 10-fold, as compared to ribavirin which reduced viral titer by 2.5 log units, or ~300-fold. Analog **5o** showed a 100-fold, or 2 log, reduction. Improvements in cell protection against *h*RSV did not translate to significant improvement in the plaque assay as was seen with ribavirin. Consequently, the titer reduction assay was not used to drive SAR efforts.

CONCLUSION

In summary, the HTS and chemistry optimization efforts produced a series of enantiomerically pure, sulfonylpyrrolidine-based compounds that are effective *in vitro* inhibitors of *h*RSV in the low micromolar range. Many of these compounds were shown to reduce the *in vitro* viral titer by 100-fold. The therapeutic index for the series was maximized at 13.4-fold and is an issue for further refinement preceding *in vivo* assessment.

EXPERIMENTAL SECTION

Chemistry

All final compounds were confirmed to be of >95% purity based on HPLC analysis. ¹H and ¹³C NMR spectra were recorded on a Bruker AM 400 spectrometer (operating at 400 and 101 MHz respectively) or a Bruker AVIII spectrometer (operating at 500 and 126 MHz respectively) in CDCl₃ with 0.03% TMS as an internal standard or DMSO-*d*₆. The chemical shifts (δ) reported are given in parts per million (ppm) and the coupling constants (*J*) are in Hertz (Hz). The spin multiplicities are reported as s = singlet, bs = broad singlet, d = doublet, t = triplet, q = quartet, dd = doublet of doublet and m = multiplet. The LCMS analysis was performed on an Agilent 1200 RRL chromatograph with photodiode array UV detection and an Agilent 6224 TOF mass spectrometer. The chromatographic method utilized the following parameters: a Waters Acquity BEH C-18 2.1 \times 50mm, 1.7 μ m column; UV detection wavelength = 214 nm; flow rate = 0.4ml/min; gradient = 5 – 100% acetonitrile over 3 minutes with a hold of 0.8 minutes at 100% acetonitrile; the aqueous mobile phase contained 0.15% ammonium hydroxide (v/v). The mass spectrometer utilized the following parameters: an Agilent multimode source which simultaneously acquires ESI+/APCI+; a reference mass solution consisting of purine and hexakis(1H, 1H, 3H-tetrafluoropropoxy) phosphazine; and a make-up solvent of 90:10:0.1 MeOH:Water:Formic Acid which was introduced to the LC flow prior to the source to assist ionization. Melting points were determined on a Stanford Research Systems OptiMelt apparatus.

Synthesis of (S)-N-(2,5-Dimethylphenyl)-1-(quinolin-8-ylsulfonyl)pyrrolidine-2-carboxamide (**5o**)

To a mixture of *L*-proline (0.50 g, 4.34 mmol) in 10% K₂CO₃ (10 mL) and THF (10 mL) was added 8-quinolinesulfonyl chloride (1.98 g, 8.68 mmol), and the resulting mixture was stirred at 50 $^{\circ}$ C for 5 h. After cooling to room temperature, the reaction mixture was acidified with 3 N aqueous HCl to pH 2 and then extracted with EtOAc (3 \times 30 mL). Separation and drying of the combined organic extracts (MgSO₄), followed by removal of solvent under reduced pressure afforded (S)-1-(quinolin-8-ylsulfonyl)pyrrolidine-2-carboxylic acid as a white solid (0.80 g, 60% yield) that did not require further purification and was used in the next step. To a solution of (S)-1-(quinolin-8-ylsulfonyl)pyrrolidine-2-

carboxylic acid (0.060 g, 0.20 mmol) in DMF (0.75 mL) was added 2,5- dimethylaniline (0.024 mL, 0.20 mmol), HATU (0.082 g, 0.22 mmol), and DIPEA (0.097 mL, 0.59 mmol). The reaction mixture was stirred for 2 h at room temperature, then diluted with CH₂Cl₂ (5 mL) and washed sequentially with aqueous 10% HCl (2 × 5 mL), saturated aqueous NaHCO₃ (2 × 5 mL), and water (2 × 5 mL). The separated organic extracts were dried (MgSO₄), and evaporated to give the crude product which was purified by silica gel flash column chromatography (2% MeOH in CH₂Cl₂) to afford (*S*)-*N*-(2,5-dimethylphenyl)-1-(quinolin-8-ylsulfonyl)pyrrolidine-2-carboxamide as a colorless oil (0.050 g, 62% yield). ¹H NMR (500 MHz; CDCl₃): δ (ppm) 9.54 (s, 1H), 8.88 (dd, *J* = 4.3 and 1.8 Hz, 1H), 8.62 (dd, *J* = 7.4 and 1.4 Hz, 1H), 8.28 (dd, *J* = 8.4 and 1.7 Hz, 1H), 8.11 (dd, *J* = 8.2 and 1.3 Hz, 1H), 7.69 (t, *J* = 7.8 Hz, 1H), 7.63 (s, 1H), 7.53 (dd, *J* = 8.3 and 4.3 Hz, 1H), 7.11 (d, *J* = 7.7 Hz, 1H), 6.93 (d, *J* = 7.6 Hz, 1H), 5.41 (dd, *J* = 7.9 and 2.0 Hz, 1H), 3.44-3.32 (m, 2H), 2.52-2.40 (m, 1H), 2.33 (s, 6H), 1.96-1.82 (m, 2H), 1.82-1.72 (m, 1H). ¹³C NMR (126 MHz; CDCl₃): δ (ppm) 170.66, 151.58, 143.91, 137.12, 136.37, 135.88, 135.29, 134.92, 134.62, 130.45, 129.29, 127.65, 126.34, 125.84, 124.23, 122.51, 63.20, 49.29, 30.19, 24.98, 21.24, 17.65. LCMS purity (214 nm) =100%. HRMS: *m/z* calcd for C₂₂H₂₃N₃O₃S (M + H⁺) 410.1533, found 410.1532. Enantiomeric excess was determined by HPLC analysis: [α]_D²⁵ −31.5 (c 0.0039 CHCl₃), > 99% ee.

Time of Addition Assay

HEp-2 cells were plated in 96 well black tissue culture plates at 10,000 cells per well in 100 μL and incubated 24 h at 37 °C, 5% CO₂. Test compounds were diluted in media to give a final concentration of 25 μM and added to plates in triplicate at −1, 0, 1, 2, 3, 5, 7, 21 and 24 h post-infection. Cells were infected with *h*RSV strain Long at an MOI of 3.0 at time point 0 and incubated for 6 days at 37 °C, 5% CO₂. Following a six day incubation period, the assay plates were equilibrated to room temperature for 30 min. An equal volume (100 μL) of Cell Titer-Glo reagent (Promega Inc.) was added to each well using a WellMate (Matrix, Hudson, NH) and the plates were incubated for an additional 10 min at room temperature. At the end of the incubation, luminescence was measured using a multi-label reader (PerkinElmer, Wellesley, MA) with an integration time of 0.1 s. Ribavirin was used as a control compound.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors appreciate the technical assistance of Xiaolin Xu. The authors gratefully acknowledge funding from the National Institutes of Health, grant number 1 R03 MH082403-01A1 to W.E.S. and partial funding in support of chemistry done at the University of Kansas Specialized Chemistry Center (NIH U54HG005031) and screening done at the Southern Research Specialized Biocontainment Screening Center (NIH U54 HG005034-01).

ABBREVIATIONS USED

<i>h</i>RSV	human respiratory syncytial virus
COPD	chronic obstructive pulmonary disease
GMP	guanosine monophosphate
GTP	guanosine triphosphate
IMPDH	inosine monophosphate dehydrogenase


RNA	ribonucleic acid
RNP	ribonucleoprotein
CPE	cytopathic effect
ATP	adenosine triphosphate
MLSMR	Molecular Libraries Small Molecule Repository
PBS	phosphate buffered saline
PI	post infection

REFERENCES

- Greenough A. Respiratory syncytial virus infection: clinical features, management, and prophylaxis. *Curr. Opin. Pulm. Med.* 2002; 8:214–217. [PubMed: 11981311]
- Langley GF, Anderson LJ. Epidemiology and Prevention of Respiratory Syncytial Virus Infections Among Infants and Young Children. *Pediatr. Infect. Dis. J.* 2011; 30:510–517. [PubMed: 21487331]
- Englund JA, Piedra PA, Whimbey E. Prevention and treatment of respiratory syncytial virus and parainfluenza viruses in immunocompromised patients. *Am. J. Med.* 1997; 102:61–70. discussion 75–76. [PubMed: 10868145]
- Jeffcoate TN. Vaccine against respiratory syncytial virus. *Lancet.* 1969; 7615:311.
- Hemming, VG. Respiratory syncytial virus: a brief history, in Contemporary diagnosis and management of respiratory syncytial virus. In: Weisman, LE.; Groothuis, JR., editors. *Handbooks in Health Care Co.* Newtown, PA: 2000. p. 7-23.
- American Academy of Pediatrics Committee on Infectious Diseases, Committee on Fetus and Newborn. Respiratory syncytial virus immune globulin intravenous: indications for use. *Pediatrics.* 1998; 99:645–650.
- Rodriguez WJ, Kim HW, Brandt CD, Fink RJ, Getson PR, Arrobio J, Murphy TM, McCarthy V, Parrott RH. Aerosolized ribavirin in the treatment of patients with respiratory syncytial virus disease. *Pediatr. Infect. Dis. J.* 1987; 6:159–163. [PubMed: 3550679]
- Smith DW, Frankel LR, Mathers LH, Tang ATS, Ariagno RL, Prober CG. A controlled trial of aerosolized ribavirin in infants receiving mechanical ventilation for severe respiratory syncytial virus infection. *New. Engl. J. Med.* 1991; 325:24–29. [PubMed: 1904551]
- Bowden RA. Respiratory virus infections after marrow transplant: the Fred Hutchinson Cancer Research Center experience. *Am. J. Med.* 1997; 102:27–30. discussion 42–43. [PubMed: 10868139]
- Leyssen P, Balzarini J, De Clercq E, Neyts J. The predominant mechanism by which ribavirin exerts its antiviral activity in vitro against flaviviruses and paramyxoviruses is mediated by inhibition of IMP dehydrogenase. *J. Virol.* 2005; 79:1943–1947. [PubMed: 15650220]
- Markland W, McQuaid TJ, Jain J, Kwong AD. Broad-spectrum antiviral activity of the IMP dehydrogenase inhibitor VX-497: a comparison with ribavirin and demonstration of antiviral additivity with alpha interferon. *Antimicrob. Agents Ch.* 2000; 44:859–866.
- Fernandez-Larsson R, O'Connell K, Koumans E, Patterson JL. Molecular analysis of the inhibitory effect of phosphorylated ribavirin on the vesicular stomatitis virus in vitro polymerase reaction. *Antimicrob. Agents Ch.* 1989; 33:1668–1673.
- Eriksson B, Helgstrand E, Johansson NG, Larsson A, Misiorny A, Noren JO, Philipson L, Stenberg K, Stening G, Stridh S, Oberg B. Inhibition of influenza virus ribonucleic acid polymerase by ribavirin triphosphate. *Antimicrob. Agents Ch.* 1977; 11:946–951.
- Gabrielsen B, Phelan MJ, Barthel-Rosa L, See C, Huggins JW, Kefauver DF, Monath TP, Ussery MA, Chmurny GN, Schubert EM, Upadhy K, Kwong C, Carter DA, Secrist JA III, Kirsi JJ, Shannon WM, Sidwell RW, Kini GD, Robins RK. Synthesis and antiviral evaluation of N-carboxamidine-substituted analogues of 1-beta-D-ribofuranosyl-1,2,4-triazole-3-carboxamidine hydrochloride. *J. Med. Chem.* 1992; 35:3231–3238. [PubMed: 1507208]

15. Douglas JL, Panis ML, Ho E, Lin K-Y, Krawczyk SH, Grant DM, Cai R, Swaminathan S, Cihlar T. Inhibition of respiratory syncytial virus fusion by the small molecule VP-14637 via specific interactions with F protein. *J. Virol.* 2003; 77:5054–5064. [PubMed: 12692208]
16. Cianci C, Yu K-L, Combrink K, Sin N, Pearce B, Wang A, Civiello R, Voss S, Luo G, Kadow K, Genovesi EV, Venables B, Gulgeze H, Trehan A, James J, Lamb L, Medina I, Roach J, Yang Z, Zadjura L, Colonna R, Clark J, Meanwell N, Krystal M. Orally active fusion inhibitor of respiratory syncytial virus. *Antimicrob. Agents Ch.* 2004; 48:413–422.
17. Ding W-D, Mitsner B, Krishnamurthy G, Aulabaugh A, Hess CD, Zaccardi J, Cutler M, Feld B, Gazumyan A, Raifeld Y, Nikitenko A, Lang SA, Gluzman Y, O'Hara B, Ellestad GA. Novel and specific respiratory syncytial virus inhibitors that target virus fusion. *J. Med. Chem.* 1998; 41:2671–2675. [PubMed: 9667956]
18. Razinkov V, Gazumyan A, Nikitenko A, Ellestad G, Krishnamurthy G. RFI-641 inhibits entry of respiratory syncytial virus via interactions with fusion protein. *Chem. Biol.* 2001; 8:645–659. [PubMed: 11451666]
19. Douglas JL. In search of a small-molecule inhibitor for respiratory syncytial virus. *Expert. Rev. Anti. Infect. Ther.* 2004; 2:625–639. [PubMed: 15482225]
20. Wyde PR, Chetty SN, Timmerman P, Gilbert BE, Andries K. Short duration aerosols of JNJ 2408068 (R170591) administered prophylactically or therapeutically protect cotton rats from experimental respiratory syncytial virus infection. *Antivir. Res.* 2003; 60:221–231. [PubMed: 14638398]
21. Sudo K, Miyazaki Y, Kojima N, Kobayashi M, Suzuki H, Shintani M, Shimizu Y. YM-53403, a unique anti-respiratory syncytial virus agent with a novel mechanism of action. *Antivir. Res.* 2005; 65:125–131. [PubMed: 15708639]
22. Liuzzi M, Mason SW, Cartier M, Lawetz C, McCollum RS, Dansereau N, Bolger G, Lapeyre N, Gaudette Y, Lagacé L, Massariol M-J, Dô F, Whitehead P, Lamarre L, Scouteau E, Bordeleau J, Landry S, Rancourt J, Fazal G, Simoneau B. Inhibitors of respiratory syncytial virus replication target cotranscriptional mRNA guanylation by viral RNA-dependent RNA polymerase. *J. Virol.* 2005; 79:13105–13115. [PubMed: 16189012]
23. Carter MC, Alber DG, Baxter RC, Bithell SK, Budworth J, Chubb A, Cockerill GS, Dowdell VCL, Henderson EA, Keegan SJ, Kelsey RD, Lockyer MJ, Stables JN, Wilson LJ, Powell KL. 1,4-Benzodiazepines as inhibitors of respiratory syncytial virus. *J. Med. Chem.* 2006; 49:2311–2319. [PubMed: 16570927]
24. Ni L, Zhao L, Qian Y, Zhu J, Jin Z, Chen YW, Tien P, Gao GF. Design and characterization of human respiratory syncytial virus entry inhibitors. *Antivir. Ther.* 2005; 10:833–840. [PubMed: 16312179]
25. Sidwell RW, Barnard DL. Respiratory syncytial virus infections: recent prospects for control. *Antivir. Res.* 2006; 71:379–390. [PubMed: 16806515]
26. Olszewska W, Ispas G, Schnoeller C, Sawant D, Van de Castele T, Nauwelaers D, Van Kerckhove B, Roymans D, De Meulder M, Rouan MC, Van Remoortere P, Bonfanti JF, Van Velsen F, Koul A, Vanstockem M, Andries K, Sowinski P, Wang B, Openshaw P, Verloes R. Antiviral and lung protective activity of a novel respiratory syncytial virus fusion inhibitor in a mouse model. *Eur. Respir. J.* 2011; 38:401–408. [PubMed: 21148224]
27. Informa UK Ltd. *Pharmaprojects – Pharmaceutical Research & Development Pipeline Intelligence.* 2008
28. Bonavia A, Franti M, Pusateri Keaney E, Kuhen K, Seepersaud M, Radetich B, Shao J, Honda A, Dewhurst J, Balabanis K, Monroe J, Wolff K, Osborne C, Lanieri L, Hoffmaster K, Amin J, Markovits J, Broome M, Skuba E, Cornella-Taracido I, Joberty G, Bouwmeester T, Hamann L, Tallarico JA, Tommasi R, Compton T, Bushell SM. Identification of broad-spectrum antiviral compounds and assessment of the druggability of their target for efficacy against respiratory syncytial virus (RSV). *Proc. Natl. Acad. Sci. U S A.* 2011; 108:6739–6744. [PubMed: 21502533]
29. Solubility and stability data was provided courtesy of the Sanford Burnham Institute.
30. Severson WE, McDowell M, Ananthan S, Chung DH, Rasmussen L, Sosa MI, White EL, Noah J, Jonsson CB. High-throughput screening of a 100,000-compound library for inhibitors of influenza A virus (H3N2). *J. Biomol. Screen.* 2008; 13:879–887. [PubMed: 18812571]

31. While compound **5b** shows inherent cellular toxicity, the observed levels of cellular viability for infected cells treated with **5b** did not appear so strongly affected such that a CPE was observed, albeit at the lower range of the three tested compounds in the series.
32. McKimm-Breschkin JL. A simplified plaque assay for respiratory syncytial virus-direct visualization of plaques without immunostaining. *J. Virol. Methods*. 2004; 120:113–117. [PubMed: 15234816]



ribavirin, 1

Figure 1.
Structure of ribavirin 1




Figure 2.
Shaded regions of hit compound 2 optimized by structure-activity relationships.











Figure 3.
Comparison of ribavirin with analogs **5b**, **5o** and **5t** in a time of addition assay.

**Scheme 1. Chemical Synthesis of Sulfonylpyrrolidine Analogs**

Reagents and conditions: (a) *L*-proline, 10% aq. K₂CO₃, THF, 50 °C, 5 h, then 3N HCl; (b) DIPEA, HATU, substituted aniline, DMF, rt, 2 h.

Table 1

hRSV CPE Assay Potency, Cytotoxicity, Selectivity Index and Logarithmic Reduction in Viral Plaques for Analogs with Structural Variation in the Proline Linker Region of the Hit Sulfonylpyrrolidine Scaffold.

entry	compound	linker structure	CPE potency \pm standard deviation (EC ₅₀ , μ M) ^a	cytotoxicity \pm standard deviation (CC ₅₀ , μ M) ^b	selectivity index (CC ₅₀ /EC ₅₀)	Virus titer log reduction (plaque assay) ^c
		linker structure				
1	1, ribavirin	NA	28.4 \pm 3.8	113.9 \pm 38.5	4.0	2.5
2	2		5.0 \pm 1.4	31.5 \pm 5.7	6.3	2.1
3	5a		> 50.0	38.1 \pm 2.3	0.8	NT
4	5b		2.7 \pm 0.9	31.8 \pm 7.7	11.8	1.0
5	5c		> 50.0	> 50.0	1.0	NT
6	5d		> 50.0	> 50.0	1.0	NT
7	5e		> 50.0	> 50.0	1.0	NT

^aData is an average of 3 experiments;

^bData is an average of 2 experiments;

\$watermark-text

\$watermark-text


\$watermark-text

^c_{NT} = not tested;

NA = not applicable; Data were analyzed using Microsoft® Excel 2010.

Table 2

hRSV CPE Assay Potency, Cytotoxicity, Selectivity Index and Logarithmic Reduction in Viral Plaques for Analogs with Structural Variation in the Aryl Amide Region of the Sulfonylpyrrolidine Scaffold.

entry	compound	 R	CPE potency \pm standard deviation (EC_{50} , μM) ^a	cytotoxicity \pm standard deviation (CC_{50} , μM) ^b	selectivity index (CC_{50}/EC_{50})	Virus titer log reduction (plaque assay) ^c
1	1 , ribavirin	NA	28.4 \pm 3.8	113.9 \pm 38.5	3.6	2.5
2	5b	2,4-dimethylphenyl	2.7 \pm 0.9	31.8 \pm 7.7	13.1	1.0
3	5f	2-methylphenyl	23.8 \pm 10.2	> 50.00	2.1	NT
4	5g	3-methylphenyl	> 50.0	25.5 \pm 4.1	0.5	NT
5	5h	4-methylphenyl	> 50.0	31.7 \pm 2.9	0.6	NT
6	5i	phenyl	> 50.0	35.6 \pm 2.3	0.7	NT
7	5j	2-ethylphenyl	21.2 \pm 5.1	> 50.0	2.6	NT
8	5k	2- <i>i</i> -propylphenyl	17.6 \pm 0.9	29.9 \pm 3.5	1.7	NT
9	5l	2,4-dimethyl-3-pyridine	> 50.0	> 50.0	1.0	NT
10	5m	3,5-dimethylphenyl	> 50.0	31.3 \pm 3.5	0.6	NT
11	5n	2,4-dichlorophenyl	> 50.0	> 50.0	1.0	NT
12	5o	2,5-dimethylphenyl	2.3 \pm 0.8	30.9 \pm 1.1	13.4	2.0
13	5p	2,6-dimethylphenyl	> 50.0	> 50.00	1.0	NT
14	5r	2-methyl-5-chlorophenyl	2.2 \pm 0.6	10.0 \pm 1.3	4.3	NT
15	5s	2-methyl,5-trifluoromethylphenyl	> 50.0	4.8 \pm 1.7	0.1	NT
16	5t	2-methyl,5-methoxyphenyl	4.2 \pm 1.3	44.2 \pm 5.7	10.4	1.0
17	5u	2-methyl,5- <i>i</i> -propylphenyl	> 50.0	9.7 \pm 1.2	0.2	NT
18	5v	2-methyl,5- <i>t</i> -butylphenyl	> 50.0	6.5 \pm 0.6	0.1	NT
19	5w	2-chloro,5-methylphenyl	> 50.0	> 50.0	1.0	NT
20	5x	2-methoxy,5-methylphenyl	> 50.0	> 50.0	1.0	NT
21	5y	(1-(<i>t</i> -butyl)-3-methyl-1H-pyrazol-5-yl)	> 50.0	> 50.0	1.0	NT

\$watermark-text

\$watermark-text

\$watermark-text

^aData is an average of 3 experiments;^bData is an average of 2 experiments;^cNT = not tested;

NA = not applicable; Data were analyzed using Microsoft® Excel 2010.